

PSEUDO-ANOSOV MAPS AND SIMPLE CLOSED CURVES ON SURFACES

SHICHENG WANG¹, YING-QING WU AND QING ZHOU¹

ABSTRACT. Suppose $\mathcal{C}$ and $\mathcal{C}'$ are two sets of simple closed curves on a hyperbolic surface F . We will give necessary and sufficient conditions for the existence of a pseudo-Anosov map g such that $g(\mathcal{C}) \cong \mathcal{C}'$.

Pseudo-Anosov maps are the most important class among surface homeomorphisms [Th3], and they play important roles in 3-manifold theory [Th2].

This note is to address the following question: Suppose F is an orientable closed surface with $\chi(F) < 0$, and suppose c and c' are two essential circles on F such that

- (1) there is an orientation preserving homeomorphism $f : F \rightarrow F$ such that $f(c) = c'$, and
- (2) c and c' are not in the same isotopy class.

Does there exists an orientation preserving pseudo-Anosov map $g : F \rightarrow F$ with $g(c) \cong c'$?

Clearly the conditions (1) and (2) are necessary for a positive answer of the question. Moreover if we replace “ F ” by “the torus”, and “pseudo-Anosov” by “Anosov” in the question, then it is known that the answer is yes. Note also if both c and c' in the Question are non-separating, then there is an orientation preserving homeomorphism f such that $f(c) = c'$.

The question is raised by M. Boileau and S. Wang to understand degree one maps between hyperbolic 3-manifolds which are surface bundles over the circle [BW]. Corollary 2 below gives an affirmative answer to this question. We need some definitions in order to state and prove our main theorem.

A circle c on a compact surface F is *essential* if c is not contractible or boundary parallel. A set $\mathcal{C} = \{c_1, \dots, c_n\}$ of mutually disjoint circles on F is an *independent*

¹ Supported by NSF of China.

set if the curves in $\mathcal{C}$ are essential and mutually non parallel. Write $\mathcal{C} \cong \mathcal{C}'$ if $\mathcal{C}$ is isotopic to $\mathcal{C}'$. An f -orbit in $\mathcal{C}$ is a nonempty subset $\mathcal{C}_1$ of $\mathcal{C}$ such that $f(\mathcal{C}_1) \cong \mathcal{C}_1$. Notice that if no curve of $\mathcal{C}$ is isotopic to a curve in $f(\mathcal{C})$, then $\mathcal{C}$ contains no f -orbit.

Theorem 1. *Let $f : F \rightarrow F$ be an orientation preserving homeomorphism on a surface F with $\chi(F) < 0$, and let $\mathcal{C}$ be an independent set of circles on F . Then there is an orientation preserving pseudo Anosov map $g : F \rightarrow F$ with $g(\mathcal{C}) \cong f(\mathcal{C})$ if and only if $\mathcal{C}$ contains no f -orbit.*

Corollary 2. *Let c and c' be non isotopic essential curves on a hyperbolic surface F such that (F, c) is orientation preserving homeomorphic to (F, c') . Then there is an orientation preserving pseudo Anosov map $g : F \rightarrow F$ such that $g(c) \cong c'$.*

Proof. This follows immediately from Theorem 1, noticing that if $\mathcal{C}$ has only one curve c then it contains an f -orbit if and only if $f(c) \cong c$. $\square$

It is not difficult to see that the condition is necessary. Below we first assume that $\mathcal{C}$ contains no f -orbit, and proceed to show that there is a pseudo-Anosov map $g : F \rightarrow F$ with $g(\mathcal{C}) \cong f(\mathcal{C})$. We remark that it is possible that $\mathcal{C}$ contains no f -orbit, and yet $f^k(c) \cong c$ for some $k > 1$, in which case we can still find a pseudo-Anosov map g such that $g(\mathcal{C}) \cong f(\mathcal{C})$.

Given a set of curves $\mathcal{C}$ on F , denote by $N(\mathcal{C})$ a regular neighborhood of $\mathcal{C}$ in F . The set $\mathcal{C}$ is a *maximal* independent set if $\mathcal{C}$ is independent and no component of $F - \text{Int}N(\mathcal{C})$ contains essential circles, that is, each component of $F - \text{Int}N(\mathcal{C})$ is a pair of pants. The following lemma allows us to replace $\mathcal{C}$ in the theorem with a maximal independent set.

Denote by τ_c a right hand Dehn twist along a circle c on F .

Lemma 3. *Let $f : F \rightarrow F$ be a homeomorphism and $\mathcal{C}$ an independent set of circles on F . If $\mathcal{C}$ contains no f -orbit, then there is a homeomorphism $f' : F \rightarrow F$ and a maximal independent set $\mathcal{C}'$ of circles such that*

- (1) $\mathcal{C} \subset \mathcal{C}'$,
- (2) $f'(c) = f(c)$ for each $c \in \mathcal{C}$, and
- (3) $\mathcal{C}'$ contains no f' -orbit.

Proof. Assume that $\mathcal{C}$ is not maximal, so there is a component S of $F - \text{Int}N(\mathcal{C})$ with $\chi(S) < 0$ which is not a disc with two holes. Now one can find a pair of essential circles c' and c'' in S such that their minimum geometric intersection number is

positive. Let $\mathcal{C}' = \mathcal{C} \cup c'$. Then $\mathcal{C}'$ is independent. Let $f_k = f \circ \tau_{c''}^k$, where k is an integer. Then for any k , $f_k(c) = f(c)$ for $c \in \mathcal{C}$. Since the minimum geometric intersection number of c' and c'' is positive, there are infinitely many isotopy classes of essential circles on F in the family $\{\tau_{c''}^k(c') \mid k \in \mathbb{Z}\}$. Since there are only finitely many curves in the set $f^{-1}(\mathcal{C}')$, we may pick k large enough so that $\tau_{c''}^k(c')$ is not isotopic to any curve in $f^{-1}(\mathcal{C}')$. Let $f' = f_k$ for this k . Then $f'(c') = f(\tau_{c''}^k(c'))$ is not isotopic to any curve in $\mathcal{C}'$.

If $\mathcal{C}'$ would contain an f' -orbit $\mathcal{C}_1$, then $f'(\mathcal{C}_1) \cong \mathcal{C}_1$, so $c' \notin \mathcal{C}_1$ because $f'(c')$ is not isotopic to any curve in $\mathcal{C}'$. But this would imply that $\mathcal{C}_1 \subset \mathcal{C}$, and since $f = f'$ on $\mathcal{C}$, it would contradict our assumption that $\mathcal{C}$ contains no f -orbit. Hence $\mathcal{C}'$ contains no f' -orbit. The proof now follows by induction because the number of curves in an independent set is bound above by $3g + h - 3$, where g is the genus and h the number of boundary components of F . $\square$

Because of Lemma 3, we may assume from now on that the independent set $\mathcal{C}$ in Theorem 1 is maximal. Let $M = F \times [0, 1]/f$ be the surface bundle over S^1 with fiber F and gluing map f , that is, it is the quotient of $F \times [0, 1]$ obtained by identifying $(x, 0)$ with $(f(x), 1)$. Let $q : F \times [0, 1] \rightarrow M$ be the quotient map. Denote by F^* the surface $q(F \times 0) = q(F \times 1)$, by $\mathcal{C}^*$ the curves $q(\mathcal{C} \times 0) = q(f(\mathcal{C}) \times 1)$ in M , and by c_i^* the curves $q(c_i \times 0) = q(f(c_i) \times 1)$ in $\mathcal{C}^*$, $i = 1, \dots, n$. Let $N(\mathcal{C}^*)$ be a tubular neighborhood of $\mathcal{C}^*$ in M ; let $M^* = M - \text{Int}N(\mathcal{C}^*)$; let T_i be the torus $\partial N(c_i^*)$ on ∂M^* .

Pick a meridian-longitude pair for each T_i , with longitude an intersection curve of $F^* \cap T_i$. Thus the slopes (i.e isotopy classes of simple closed curves) on T_i are in one to one correspondence with the numbers in $\mathbb{Q} \cup \{\infty\}$. (See [R] for more details.) Define $M^*(q_1, \dots, q_n)$ to be the manifold obtained by q_i Dehn filling on T_i , $i = 1, \dots, n$.

Lemma 4. *Let $g(k_1, \dots, k_n) = f \circ \tau_{c_1}^{k_1} \circ \dots \circ \tau_{c_n}^{k_n}$. Then*

- (1) $g(k_1, \dots, k_n)(c_i) = f(c_i)$ for all $i = 1, \dots, n$ and $k_j \in \mathbb{Z}$.
- (2) $F \times [0, 1]/g(k_1, \dots, k_n) = M^*(1/k_1, \dots, 1/k_n)$.

Proof. (1) Since the curves in $\mathcal{C}$ are mutually disjoint, $\tau_{c_j}(c_i) = c_i$ for all i, j ; hence $g(k_1, \dots, k_n)(c_i) = f \circ \tau_{c_1}^{k_1} \circ \dots \circ \tau_{c_n}^{k_n}(c_i) = f(c_i)$.

(2) This is a simple geometric observation: Twisting along the curves will not change M^* , the exterior of the link $\mathcal{C}^*$ in M , while a meridian of c_i^* in $F \times [0, 1]/f$

is changed to a $1/k_i$ curve in $F \times [0, 1]/g(k_1, \dots, k_n)$. $\square$

We would like to show that $g(k_1, \dots, k_n)$ are isotopic to pseudo-Anosov maps for some k_i . By a deep theorem of Thurston (see Otal [Ot] for proof), it suffices to show that $M^*(1/k_1, \dots, 1/k_n)$ is hyperbolic. It is attempting to apply the hyperbolic surgery theorem of Thurston [Th1], but we have to be careful because the manifold M^* may not be hyperbolic.

By an isotopy we may assume that either $f(c_i) = c_j$ or $f(c_i)$ is not isotopic to c_j . Define an oriented graph Γ associated to $(\mathcal{C}, f)$ as follows. Each element c_i in $\mathcal{C}$ is a vertex of Γ , still denoted by c_i , and there is an oriented edge $c_i c_j$ in Γ if and only if $c_j = f(c_i)$. A subgraph of Γ is a *chain* if it is homeomorphic to an interval. Since there is no f -orbit and the curves $f(c_i)$ are mutually non isotopic, we have

Lemma 5. *Γ has only finitely many components, each of which is either a single vertex or a chain.*

For each component γ of Γ , define C_γ in M as follows: If γ is an isolated vertex c_i , let C_γ be the knot c_i^* . If γ has an oriented edge $c_i c_j$, let $C_{c_i c_j}$ be the annulus $q(c_j \times [0, 1])$ in M . Notice that since $f(c_i) = c_j$, we have

$$q(c_j \times 1) = q(f(c_i) \times 1) = q(c_i \times 0) = c_i^*.$$

Thus $C_{c_i c_j}$ is an annulus with “top” boundary curve c_i^* and “bottom” boundary curve c_j^* . Now define C_γ to be the union of $C_{c_i c_j}$ over all edges $c_i c_j$ of γ . Then C_γ is an embedded annulus in M containing all $\{c_i^* \mid c_i \in \gamma\}$. Different components of Γ correspond to disjoint C_γ in M . Let C_Γ be the union of C_γ for all $\gamma \subset \Gamma$.

Lemma 6. *Every incompressible torus in M^* is isotopic to a torus in $N(C_\Gamma)$.*

Proof. Since F^* is incompressible in the surface bundle M , the surface $F' = F^* \cap M^*$ is also incompressible in M^* . Let T be an incompressible torus in M^* . We assume that T has been isotoped in M^* to meet F' minimally, so $\beta = T \cap F'$ consists of essential curves on both T and F' ; in particular, it cuts T into π_1 injective annuli in M^* . Note that $\beta \neq \emptyset$, otherwise T would be an incompressible torus in $F \times [0, 1]$; which is impossible as $\chi(F) < 0$.

Since $\mathcal{C}$ is maximal, each component of F' is a disk with two holes, hence each component of β is parallel, on F^* , to a unique curve c_i^* in $\mathcal{C}$. We may isotope T so that β lies in $N(C_\Gamma)$.

An annulus A in $F \times [0, 1]$ is called a *vertical* annulus if it is isotopic to a product $s \times [0, 1]$ for a simple closed curve s in F ; it is a *horizontal* annulus if it is boundary parallel. Each π_1 -injective annulus in $F \times [0, 1]$ is either horizontal or vertical.

Cutting M^* along F' , we get a manifold M' homeomorphic to $F \times [0, 1]$. If a component A_k of $T \cap M'$ is horizontal then its two boundary components are isotopic to the same c_i^* on F^* . Since $\partial A_k \subset \beta$ is already in $N(C_\Gamma)$, by an isotopy rel ∂A_k we may push A_k into $N(c_i^*) \subset N(C_\Gamma)$. If it is vertical, let $c_i^* = c_i \times 0$ and $c_j^* = f(c_j) \times 1$ be the two curves on $F \times \partial[0, 1]$ isotopic to ∂A_k . Then the annulus A_k gives rise to an isotopy between c_i and $f(c_j)$ on F , so by our assumption above, we have $c_i = f(c_j)$, so $C_{c_i c_j} \subset C_\Gamma$. It follows that A_j is also rel ∂A_j isotopic into $N(C_\Gamma)$. This completes the proof of Lemma 6. $\square$

Lemma 7. $X = M - \text{Int}N(C_\Gamma)$ is a hyperbolic manifold.

Proof. X is irreducible: A reducing sphere S would bound a ball B in M because M , as an F bundle over S^1 , is irreducible. Hence B contains some component of C_Γ . But since each component of C_Γ contains some essential curve c_i^* of F , and since F is π_1 injective in M , this is impossible.

X is not a Seifert fiber space: Each component of $N(C_\Gamma)$ can be shrunk to some $N(c_i^*)$, so X contains a nonseparating, hyperbolic, closed incompressible surface, i.e., the image of $F \times \frac{1}{2}$ under the reverse isotopy of the above shrinking process. No such surface exists in a Seifert fiber space with boundary because an essential surface in such a manifold is either horizontal (hence bounded) or vertical (hence a torus).

X is also atoroidal: If T is an essential torus in X , then by Lemma 6 there is a torus T' in some $N(C_\gamma) \cap M^*$ such that $T \cup T'$ bounds a product region $W = T \times [0, 1]$ in M^* . Since T is essential, T' is π_1 injective in $N(C_\gamma) \cap M^* = P \times S^1$, where P is a planar surface. Note that a π_1 injective torus in $P \times S^1$ is isotopic to $c \times S^1$ for some circle c in P . Since $W \subset M^*$, it contains no component of ∂M^* , so c must be parallel to the boundary curve of P on $\partial N(C_\gamma)$, which means that T' , hence T , is parallel to $\partial N(C_\gamma)$.

Since ∂M is a union of tori, the lemma now follows from the Geometrization Theorem of Thurston for Haken manifolds [Th2]. $\square$

Proof of Theorem 1. If $\mathcal{C}$ contains an f -orbit $\mathcal{C}_1$, then there is a circle $c \in \mathcal{C}$ such that $f^i(c)$ is isotopic to a curve in $\mathcal{C}_1$ for all i , and $f^k(c) \cong c$ for some $k \neq 0$.

Suppose $g : F \rightarrow F$ is a homeomorphism such that $g(c') \cong f(c')$ for all $c' \in \mathcal{C}$, then by induction we have

$$g^i(c) = g(g^{i-1}(c)) \cong g(f^{i-1}(c)) \cong f^i(c)$$

for all i . In particular, $g^k(c) \cong c$, so g cannot be a pseudo-Anosov map.

Now suppose $\mathcal{C}$ contains no f -orbit. Let $\gamma_1, \dots, \gamma_m$ be the distinct components of Γ . After relabeling, we may assume that c_i is a vertex of γ_i for $i \leq m$. Since $N(C_{\gamma_i})$ is isotopic to $N(c_i^*)$, by performing trivial surgery on c_j^* for all $j > m$ we get a 3-manifold

$$X = M^*(\emptyset, \dots, \emptyset, \infty, \dots, \infty) = M - \text{Int}N(C_\Gamma).$$

By Lemma 7, X is a hyperbolic manifold, therefore, by the Hyperbolic Surgery Theorem of Thurston [Th1], $X(1/k_1, \dots, 1/k_m)$ is hyperbolic for sufficiently large k_i . By Lemma 4(2) we have

$$X\left(\frac{1}{k_1}, \dots, \frac{1}{k_m}\right) = M^*\left(\frac{1}{k_1}, \dots, \frac{1}{k_m}, \infty, \dots, \infty\right) = F \times [0, 1]/g(k_1, \dots, k_m, 0, \dots, 0).$$

By Lemma 4(1) we have $g(k_1, \dots, k_m, 0, \dots, 0)(\mathcal{C}) = f(\mathcal{C})$. The theorem now follows from Thurston's theorem that $F \times [0, 1]/g$ is hyperbolic if and only if g is isotopic to a pseudo Anosov map [Th2, Ot]. $\square$

ACKNOWLEDGEMENT. We would like to thank R.D. Edwards, T. Kobayoshi and W. Thurston for some helpful conversations.

REFERENCES

- [BW] M. Boileau and S.C. Wang, *Non-zero degree maps and surface bundles over S^1* , J. Diff. Geom. **43** (1996), 789-806.
- [Ot] J.P. Otal, *Hyperbolisation des 3-variétés fibrés*, Asterisque. Soc. Mat. de France **110** (1997).
- [Rf] D. Rolfsen, *Knots and Links*, Publish or Perish, 1976.
- [Th1] W. Thurston, *The Geometry and Topology of 3-manifolds*, Princeton University, 1978.
- [Th2] ———, *Three dimensional manifolds, Kleinian groups and hyperbolic geometry*, Bull. Amer. Math. Soc. **6** (1982), 357–381.
- [Th3] ———, *On the geometry and dynamics of diffeomorphism of surfaces*, Bull. Amer. Math. Soc. **19** (1988), 417–431.

PEKING UNIVERSITY, BEIJING 100871, CHINA
E-mail address: swang@sxx0.math.pku.edu.cn

UNIVERSITY OF IOWA, IOWA CITY, IA 52242, USA
E-mail address: wu@math.uiowa.edu

EAST CHINA NORMAL UNIVERSITY, SHANGHAI 200062, CHINA
E-mail address: qzhou@math.ecnu.edu.cn